Modification from Noncommutative Inflation

Hong-Guang Zhang[1],* Jie Ren[2],† and Xin-He Meng[1]‡
[1] *Department of Physics, Nankai University, Tianjin 300071, China and*
[2] *Theoretical Physics Division, Chern Institute of Mathematics, Nankai University, Tianjin 300071, China*
(Dated: June 18, 2009)

We explore the noncommutative effect on single field inflation and compare with WMAP five-year data. First, we calculate the noncommutative effect from the potential and dynamical terms, and construct the general form of modified power spectrum. Second, we consider the leading order modification of slow-roll, DBI and K-inflation and unite the modification, which means the modification is nearly model independent at this level. Finally, comparing with the WMAP5 data, we find that the modified can be well realized as the origin of the relative large spectral index and the quite small running.

PACS numbers: 98.80.-k

arXiv:0801.4142v3 [astro-ph] 18 Jun 2009

I. INTRODUCTION

Inflation [1] is a very successful paradigm for naturally understanding the puzzling aspects of hot big bang cosmology such as flatness, homogeneity and monople problems. As it causally explains the origin of the super-horizon density perturbations, inflation seeds the large structure of the universe we observe today. More like a paradigm than a theory because of the missing fundamental basis, inflation is well established by the observation of cosmic microwave background. Models based on pure de sitter spacetime predict a scale-invariant and adiabatic power spectrum, which can be characterized by $n_s - 1 = 0$. However, this is not the history of our unverse. WMAP and other astrophysical observations predict nontrivial spectral index and running.

The WMAP team has reported the five-year data[2] and used them to constrain the physics of inflation. For the ΛCDM model, WMAP5 data show that the index of power spectrum satisfies $n_s = 0.963^{+0.014+0.029}_{-0.015-0.028}$ $(1\sigma, 2\sigma$ CL);. Combining WMAP with SDSS and SNIa, the result is $n_s = 0.960^{+0.014+0.026}_{-0.013-0.027}$ $(1\sigma, 2\sigma$ CL). Though the red power spectrum is still favored at the level of 3σ CL, the result is a little bluer than that of WMAP3. And, the running of the spectral index is not favored anymore. With WMAP5 data only, the running is $\alpha_s = \frac{dn_s}{d\ln k} = -0.037^{+0.028}_{-0.028}$ $(1\sigma$ CL). Combining with SDSS and SNIa data, the result is $\alpha_s = \frac{dn_s}{d\ln k} = -0.032^{+0.021}_{-0.020}$ $(1\sigma$ CL). We find that the spectral index in WMAP5 is slightly larger than that in WMAP3 and the running is much smaller than that in WMAP3.

Theoretically, we can modify the inflation models, but it will increase the implicity from the aspect of a fundamental theory. In the series works of [3, 4, 5], running index emerges from non-Bunch-Davis vacuum and the new physics will imprint on CMB. However, the serious problem is we still know little about de sitter spacetime, which is noted as the ambiguity of vacuum selection.

In the papers [6], another method has been presented. The authors obtained the nontrivial spectral index and its running from the generalized slow-roll inflation with arbitrary sound speed. However, as a new challenge for most inflation models, we expect the spectral index and its running can provide the opportunity to observe the short distance physics and the first moment scenario of our universe.

As a candidate to probe the quantum gravity in string theory, noncommutativity is applied to describe D-brane physics. Considering the string effect and strong string interaction, short distance geometry becomes very different. In addition, while the usual concept of geometry is completely break down near the singularity geometry, noncommutativity is also a good description.

In the works of [7, 8], the authors introduced noncommutativity, which naturally emerges from string theory, to inflation physics. As the running of spectral index is large, they obtained nice results with WMAP3 experiment data. However, most recent, WMAP5 explores a quite small running index.

In this lecture, we use another method to calculate the modification. We directly calculate the modification of power spectrum from the noncommutative potential and dynamical terms while the quantum modes solution still

*Electronic address: zhglost@hotmail.com
†Electronic address: jrenphysics@hotmail.com
‡Electronic address: xhm@nankai.edu.cn

remains classical. We study the modification of the single field inflation models including slow-roll, DBI and K-inflation. As our calculation includes all the potential and dynamics terms, the modification is exact. We find these three modification can be written in one unite form at leading order. This means that the modification can not be tested by different models. In addition, the leading order of modified power spectrum is $1/k^2$, which means that the modification is small. Our calculation shows that the noncommutative potential and dynamical terms can be realized as the possible origin of the spectral index and its running since it is favored by WMAP5 data.

This paper is organized as follows. In section 2, we review the inflation formalism with a general lagrangian. In section 3, we study the noncommutative effect on inflation including the modification of potential and dynamic and reconstruct the power spectrum. In section 4, we study concrete models and compare with the WMAP5 observation. Section 5 contains some conclusion and discussion.

II. FORMALISM WITH GENERAL LAGRANGIAN

In this section, we review the formalism presented in [9, 10] which can describe different inflation models in a general Lagrangian $P(X,\phi)$. The Lagrangian is of the general form

$$S=\frac{1}{2}\int d^4x\sqrt{-g}[R+2P(X,\phi)], \tag{1}$$

where ϕ is the inflation field and $X=-\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi$ is the dynamic term. We have set $M_{pl}=(8\pi G)^{-\frac{1}{2}}=1$ and the signature of the metric is $(-1,1,1,1)$. The energy of the inflaton field is defined as

$$E=2XP_{,X}-P, \tag{2}$$

where $P_{,X}$ denote the derivative with respect to X. We study the universe that is homogeneous with a Friedmann-Robertson-Walker metric

$$ds^2=-dt^2+a^2(t)dx_3^2, \tag{3}$$

where $a(t)$ is the scale factor and $H=\dot{a}/a$ is the Hubble parameter. It is useful to define the "speed of sound" c_s as

$$c_s^2=\frac{dP}{dE}=\frac{P_{,X}}{P_{,X}+2XP_{,XX}} \tag{4}$$

.

The primordial power spectrum P_k^ζ evaluated at the time of horizon exit at $c_sk=aH$ and the spectral index n_s can be derived as

$$\begin{aligned} P_k^\zeta&=\frac{1}{36\pi^2}\frac{E^2}{c_s(P+E)}=\frac{1}{8\pi^2}\frac{H^2}{c_s\epsilon}\ ,\\ n_s-1&=\frac{d\ln P_k^\zeta}{d\ln k}=-2\epsilon-\eta-s, \end{aligned} \tag{5}$$

The motive equation of inflaton is

$$(a\varphi)''+(c_s^2k^2-\frac{a''}{a})a\varphi=0. \tag{6}$$

The classical solution is

$$\varphi=\frac{\dot{\phi}_0}{\sqrt{4\epsilon c_sk^3}}(1+ic_sk\eta)e^{-ic_sk\eta}, \tag{7}$$

where we choose the standard Bunch-Davies vacuum as the background to fix the coefficient. In this paper, we study the modification of the noncommutativity effects on the potential and dynamic terms while the solution of inflaton still remains classical.

III. MODIFICATION PROCEDURE

Noncommutative geometry on small scale is a consequence of string theory. A better description of noncommutativity geometry, in terms of the algebra generated by noncommutative coordinate, is $[x^\mu, x^\nu] = i\theta^{\mu\nu}$, where $\theta^{\mu\nu}$depends on the background flux in string theory. As has been showed in paper[11], a simple form of $\theta^{\mu\nu}$ is the string length.The efficient way to define product is by so-called Moyal product, whose multiple expansion in curved space-time can be expressed as

$$f_1 \star \cdots \star f_n = (1 + \frac{i}{2}\theta^{\mu\nu}\sum_{a<b} D^a_\mu D^b_\nu - \frac{1}{8}\theta^{\mu\nu}\theta^{\rho\sigma}\sum_{a<b,c<d} D^a_\mu D^b_\nu D^c_\rho D^d_\sigma) f_1 \cdots f_n. \tag{8}$$

The lowest order noncommutative modification term is of order $O(\theta^2)$ and the lowest order of the perturbation is

$$\delta_\theta f_1 \star \cdots \star f_n = -\frac{1}{8}\theta^{\mu\nu}\theta^{\rho\sigma}\sum_{a<b,c<d} (D^a_\mu D^b_\nu D^c_\rho D^d_\sigma) f_1 \cdots f_n. \tag{9}$$

As discussed in [12], we set the spacetime component of noncommutativity to zero: $\theta^{0i} = 0$ without losing generality and considering the unitary. Moreover, considering the uncertain relation in string theory and noncommutativity in brane world sheet, we finally obtain $\theta^{12} = l_s^2/a^2$ in the comoving coordinate. We consider the correction of the lowest order θ^2 in this paper, and denote them as $\delta_\theta X$ and $\delta_\theta V$ respectively. The change of the inflation action is

$$S = \int d^4x\sqrt{-g}(P_{,X}\delta_\theta X + P_{,V}\delta_\theta V). \tag{10}$$

We divide the inflaton ϕ into the isotropic background $\phi_0(t)$ and the fluctuation φ, $\phi = \phi_0 + \varphi$. They are constraint by a new parameter ζ,

$$\zeta = \frac{H}{\dot\phi_0}\varphi + \mathcal{O}(\epsilon,\eta)(\frac{H}{\dot\phi_0}\varphi)^2, \tag{11}$$

where ϵ, η are the slow roll parameters. Considering the slow-roll constraint from experiment, we choose the gauge transformation $\zeta = \frac{H}{\dot\phi_0}\varphi$. After serially taylor expand $P_{,X}$ and $P_{,V}$ around the background value and multiply with the according terms in $\delta_\theta X$ and $\delta_\theta V$, we obtain the whole change of second order action of perturbation due to noncommutative geometry. In leading order of slow-roll parameter it can be written as

$$\begin{aligned}\delta_\theta S_2 \;=\; \int d^4x\sqrt{h}(&P_{,X_0}(\delta_\theta X)_2 + P_{,X_0X_0}(\delta_g X)_1(\delta_\theta X)_1 \\ &+P_{,V_0}(\delta_\theta V)_2 + P_{,V_0X_0}(\delta_g X)_1(\delta_\theta V)_1) \\ &+P_{,X_0\phi_0}\varphi(\delta_\theta X)_1,\end{aligned} \tag{12}$$

where $\delta_g X = X - X_0$, $(\delta_g X)_1 = \dot\phi_0\dot\varphi$, $(\delta_g X)_2 = \frac{1}{2}(\dot\varphi^2 - (\partial\varphi)^2)$ and we have picked out the terms with least $\dot\phi_0$ to reduce the order of slow-roll parameter. To simplify the calculation, we need to pick out the terms of leading order of slow-roll parameter. We decompose $\delta_\theta X$ and $\delta_\theta V$ into terms of different order of perturbation

$$\begin{aligned}\delta_\theta X &= (\delta_\theta X)_2 + (\delta_\theta X)_1 + (\delta_\theta X)_0 \\ \delta_\theta V &= (\delta_\theta V)_2 + (\delta_\theta V)_1 + (\delta_\theta V)_0.\end{aligned} \tag{13}$$

Without losing generality, we presume the potential term as $V(\phi) = \phi^n$ and the dynamic term as $X = -\frac{1}{2}\partial_\mu\phi\partial^\mu\phi$ in this paper. The authors in paper[11] have calculated $\delta_\theta X$ and $\delta_\theta V$ and their orders as A1, A2, A3.

Using the "in-in" formulism, the two-point function is calculated through

$$\langle\zeta^2(t)\rangle = -i\int_{t_0}^{t} dt'\langle[\zeta^2(t), H_{int}(t')]\rangle. \tag{14}$$

We evaluate the modification of two-point function, which is denoted by $\delta\langle\zeta^2(t)\rangle_\theta$ below, in the lowest order of θ as

$$\delta\langle\zeta(x_1)\zeta(x_2)\rangle_\theta = -2\,\mathcal{R}e(\int_{\eta_0}^{\eta} d\eta' i\langle\zeta(x_1)\zeta(x_2)(\delta L_{int} + 2(\delta_\theta V)_2)(\eta')\rangle_0). \tag{15}$$

By setting $k_1 = k_2 = k$, the explicit power spectrum modified by noncommutative correction of second order action is

$$\begin{aligned}\delta P_\theta^\zeta &= \delta\langle\zeta(x_1)\zeta(x_2)\rangle_\theta \\ &= \frac{\pi^2 l_s^4 \phi_0^2 \dot{\phi}_0^2 H}{32\epsilon^2}\Bigg[(\frac{-1}{k^2} - \frac{45}{8k^6} + \frac{681}{16c_s^4 k^{10}} - \frac{333}{8c_s^6 k^{14}})P_{,X} \\ &\quad +(\frac{c_s}{2k^2} + \frac{5\dot{\phi}_0^2}{8k^6})P_{,XX} - \frac{n(n-1)\dot{\phi}_0^{n-1}}{4H_2 k^2}P_{,V} \\ &\quad +\frac{n(n-1)\dot{\phi}_0^{n-2}}{4H^2 k^2}P_{,VX}\Bigg]. \end{aligned} \tag{16}$$

IV. MODIFICATION OF INFLATION MODELS AND COMPARISON WITH WMAP5

First, we study the Slow-roll inflation model which is the most popular model studied in the literature. The effective action takes the canonical non-relativistic form:

$$P(X,\phi) = X - V(\phi). \tag{17}$$

The value of speed of sound $c_s = 1$. The modified parts can be expressed as

$$P_{,X} = 1, \quad P_{,XX} = 0, \quad P_{,V} = -1, \quad P_{,XV} = 0. \tag{18}$$

So the modified primordial power spectrum is

$$\delta P_\theta^\zeta = \frac{\pi^2 l_s^4 \phi_0^2 \dot{\phi}^2 H}{32\epsilon^2}\left[\frac{-1}{k^2} + \frac{n(n-1)\dot{\phi}_0^{n-1}}{4H_2 k^2} - \frac{45}{8k^6} + \frac{681}{16c_s^4 k^{10}} - \frac{333}{8c_s^6 k^{14}}\right], \tag{19}$$

while the power spectrum of the slow-roll inflation without modification is

$$P_0^\zeta = C_1 k^{-2\varepsilon}, \tag{20}$$

where C_1 is a constant.

Second, we consider DBI inflation[13] which is motivated by brane inflation scenario in warped compactifications. The effective Lagrangian is

$$P(X,\phi) = -f(\phi)^{-1}\sqrt{1-2f(\phi)X} + f(\phi)^{-1} - V(\phi), \tag{21}$$

where f is the warp factor $f = \frac{\lambda}{\phi^4}$, and λ depends on flux number. The value of speed of sound is $c_s = \sqrt{1-\dot{\phi}^2 f(\phi)}$. The modified parts can be expressed as

$$P_{,X} = \frac{1}{c_s}, \quad P_{,XX} = \frac{P_{,X}}{2X_0}(\frac{1}{c_s^2} - 1), \quad P_{,V} = P_{,XV} = 0. \tag{22}$$

We find that the modification of DBI inflation is independent of the potential term. The modified parts of primordial power spectrum is

$$\begin{aligned}\delta P_\theta^\zeta &= \frac{\pi^2 l_s^4 \phi_0^2 \dot{\phi}_0^2 H}{32\epsilon^2}\Bigg[\frac{-1}{c_s k^2} + \frac{1}{4c_s X_0 k^2} + \frac{c_s}{4X_0 k^2} - \frac{45}{8c_s k^6} \\ &\quad +\frac{5\dot{\phi}_0^2}{16X_0 c_s^2 k^6} - \frac{5\dot{\phi}_0^2}{16X_0 k^6} + \frac{681}{16c_s^5 k^{10}} - \frac{333}{8c_s^7 k^{14}}\Bigg]. \end{aligned} \tag{23}$$

As presented in the paper of [14], the power spectrum of the DBI inflation is

$$P_0^\zeta = C_2 k^{-\frac{2\epsilon+s}{1-\epsilon-s}}, \tag{24}$$

where C_2 is a constant and flow parameters ϵ, s are also constant.



FIG. 1: The relation between the spectral index n_s and the comoving wave number k.



FIG. 2: The relation between the running spectral index α and the comoving wave number k.

Finally, let we consider the correction in K-inflation [10] model with small speed of sound. The Lagrangian of the power law K-inflation is of the form

$$P(X,\phi) = \frac{4}{9}\frac{4-3\gamma}{\gamma^2}\frac{1}{\phi^2}(-X+X^2), \tag{25}$$

where γ is a constant. Considering $a \sim t^{\frac{2}{3\gamma}}$, we set $0 < \gamma < \frac{2}{3}$. So the speed of sound is $c_s^2 = \frac{\gamma}{8-3\gamma}$. In order to get small speed of sound, we focus on the region $\gamma \ll 1$. The modified parts can be expressed as

$$P_{,X} = \frac{16-12\gamma}{9\gamma^2\phi^2}(2X-1), \quad P_{,XX} = 2\frac{16-12\gamma}{9\gamma^2\phi^2}, \quad P_{,V} = P_{,XV} = 0. \tag{26}$$

So, the modified primordial power spectrum of K-inflation is

$$\begin{aligned}\delta P_\theta^\zeta &= \frac{\pi^2 l_s^4 \phi_0^2 \dot{\phi_0^2} H}{32\epsilon^2}\Bigg[(\frac{-1}{k^2} - \frac{45}{8k^6} + \frac{681}{16c_s^4 k^{10}} - \frac{333}{8c_s^6 k^{14}})P_{,X} \\ &\quad +(\frac{5\dot{\phi_0^2}}{8k^6} + \frac{c_s}{2k^2})P_{,XX}\Bigg]\end{aligned} \tag{27}$$

The power spectrum of the K-inflation in the limit of small γ is

$$P_k^\zeta = C_3 k^{-3\gamma} \tag{28}$$

where C_3 is a constant. According to the above results, if we only consider the leading order corrections in terms of the comoving wave number k by noncommutative effects, the modified primordial power spectrum can be written as

$$P_k^\zeta = A(k^{-2\epsilon} - \frac{C}{k^2}), \tag{29}$$

where A, ϵ, and C are all parameters. So, the modification from noncommutativity is model independent at leading order.

The scalar spectral index derived from the WMAP5 data is relatively blue comparing to that of WMAP3, though it is still red-tilted. The running of the spectral index, according to the WMAP5 data, explores an upward shift from the three-year result, $\alpha_s = -0.032^{+0.021}_{-0.020}$ (WMAP5 + BAO + SN). Fig. 1 plots the scalar spectral index n_s as a function of the comoving wave number k. The Fig. 2 plots the running of the scalar spectral index α as a function of k.

V. CONCLUSION

As an important method to detect short distance physics, noncommutativity naturally emerges from string theory and is applied to inflation physics. Branderberger and Ho found noncommutativity can modify the power spectrum in a significant way. In addition, in papers[15], it has been used to regulate the eternal inflation.

Recent released WMAP5 data favor a red-tilted power spectrum of primordial fluctuations at the level of two standard deviations, which is the same as the WMAP3 result. However, qualitatively, the spectral index is slightly greater and the running is quite small than the three-year value.

Our calculation is quite different. First, we calculated the modification from the potential and dynamical terms while the solution of inflaton remains classic. Second, we calculated the modification of slow-roll, DBI and K-inflation. We found the modification is model independent, as we only considering the leading order. In addition, as the leading order modification of power spectrum is proportional to $1/k^2$, the modification of is quite small and can be favored by WMAP5. Third, as we used the string scale in our paper and the correction is proportional to $(\theta^{12})^2$, it could be larger if we have a relative low noncommutative scale.

Finally, as WMAP five-year data provide stringent limits on deviation from the minimal, 6-parameter ΛCDM model, we expect more years WMAP and Planck data give us more information about the anisotropy of CMB. In addition, future CMB missions such as ground-based polarization experiment of BICEP and ESA's Planck Surveyor, are also expected to help us faithfully understand the early universe.

ACKNOWLEDGEMENTS

Zhang would like to thank W. Xue, E. Lim and K. Fang for great help. We are grateful to KITPC where this work was initiated. Xin-He Meng is supported by NSFC under No. 10675062.

APPENDIX A: NONCOMMUTATIVE CORRECTION

$$\begin{aligned}
\delta_\theta X &= \frac{1}{8}(\theta^{12})^2 g^{\mu\nu}(D_1 D_1 D_\mu \phi D_2 D_2 D_\nu \phi - D_1 D_2 D_\mu \phi D_2 D_1 D_\nu \phi), \\
\delta_\theta V &= -\frac{n(n-1)}{8}(\theta^{12})^2 \phi^{n-2}(D_1 D_1 \phi D_2 D_2 \phi - D_1 D_2 \phi D_2 D_1 \phi) \\
&\quad -\frac{n(n-1)(n-2)}{24}(\theta^{12})^2 \phi^{n-3}(D_1 D_1 \phi D_2 \phi D_2 \phi + D_2 D_2 \phi D_1 \phi D_1 \phi \\
&\quad -D_1 D_2 \phi D_1 \phi D_2 \phi - D_2 D_1 \phi D_2 \phi D_1 \phi).
\end{aligned} \tag{A1}$$

$$
\begin{aligned}
(\delta_\theta V)_1 &= -\frac{n(n-1)}{8}(\theta^{12})^2\phi_0^{n-2}(2(a\dot{a})^2\dot{\phi}_0\dot{\varphi} - a\dot{a}\dot{\phi}_0\partial_1^2\varphi - a\dot{a}\dot{\phi}_0\partial_2^2\varphi).\\
(\delta_\theta V)_2 &= -\frac{n(n-1)}{8}(\theta^{12})^2\phi_0^{n-2}(\partial_1^2\varphi\partial_2^2\varphi - \partial_1\partial_2\varphi\partial_1\partial_2\varphi + (a\dot{a})^2\dot{\varphi}^2\\
&\quad -a\dot{a}\dot{\varphi}\partial_1^2\varphi - a\dot{a}\dot{\varphi}\partial_2^2\varphi).
\end{aligned}
\tag{A2}
$$

$$
\begin{aligned}
(\delta_\theta X)_1 &= -\frac{1}{8}(\theta^{12})^2 Ha\dot{a}\dot{\phi}_0(\partial_1^2\dot{\varphi} + \partial_2^2\dot{\varphi} - 2H\partial_1^2\varphi - 2H\partial_2^2\varphi + 2Ha\dot{a}\dot{\varphi}).\\
(\delta_\theta X)_2 &= \frac{1}{8}(\theta^{12})^2(\frac{1}{a^2}\partial_1^2\partial_i\varphi\partial_2^2\partial_i\varphi - \frac{1}{a^2}\partial_1\partial_2\partial_i\varphi\partial_1\partial_2\partial_i\varphi - H\partial_1^2\partial_i\varphi\partial_i\dot{\varphi}\\
&\quad -H\partial_2^2\partial_i\varphi\partial_i\dot{\varphi} + H^2\partial_1^2\partial_i\varphi\partial_i\varphi + H^2\partial_2^2\partial_i\varphi\partial_i\varphi + \dot{a}^2\partial_1\dot{\varphi}\partial_1\dot{\varphi}\\
&\quad +\dot{a}^2\partial_2\dot{\varphi}\partial_2\dot{\varphi} + \dot{a}^2\partial_i\dot{\varphi}\partial_i\dot{\varphi} - 2H\dot{a}^2\partial_1\dot{\varphi}\partial_1\varphi - 2H\dot{a}^2\partial_2\dot{\varphi}\partial_2\varphi\\
&\quad -2H\dot{a}^2\partial_i\dot{\varphi}\partial_i\varphi + (H\dot{a})^2\partial_1\varphi\partial_1\varphi + (H\dot{a})^2\partial_2\varphi\partial_2\varphi\\
&\quad +(H\dot{a})^2\partial_i\varphi\partial_i\varphi - \partial_1^2\dot{\varphi}\partial_2^2\dot{\varphi} + \partial_1\partial_2\dot{\varphi}\partial_1\partial_2\dot{\varphi} - (Ha\dot{a})^2\dot{\varphi}^2\\
&\quad -Ha\dot{a}\dot{\varphi}\partial_1^2\dot{\varphi} + Ha\dot{a}\dot{\varphi}\partial_2^2\dot{\varphi} + 2H\partial_1^2\dot{\varphi}\partial_2^2\varphi + 2H\partial_1^2\varphi\partial_2^2\dot{\varphi}\\
&\quad +2H^2a\dot{a}\dot{\varphi}\partial_1^2\varphi + 2H^2a\dot{a}\dot{\varphi}\partial_2^2\varphi - 4H\partial_1\partial_2\dot{\varphi}\partial_1\partial_2\varphi\\
&\quad -4H^2\partial_1^2\varphi\partial_2^2\varphi + 4H^2\partial_1\partial_2\varphi\partial_1\partial_2\varphi + a\dot{a}{\partial_1}^2\dot{\varphi}\ddot{\varphi} + a\dot{a}{\partial_2}^2\dot{\varphi}\ddot{\varphi}\\
&\quad -a^2\dot{a}^2\ddot{\varphi}\ddot{\varphi} - \dot{a}^2\ddot{\varphi}(2{\partial_1}^2\varphi + 2{\partial_2}^2\varphi - a\dot{a}\dot{\varphi})).
\end{aligned}
\tag{A3}
$$

[1] A. H. Guth, Phys.Rev.D 23(1981)347; A. D. Linde, Phys.Lett.B 108(1982)389 ; A.Albrecht and P. J. Steinhardt, Phys. Rev. Lett. 48(1982)220.

[2] E. Komatsu et al., Five-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation, arXiv:0803.0547.

[3] Robert H. Brandenberger, On Signatures of Short Distance Physics in the Cosmic Microwave Background, Int.J.Mod.Phys.A 17(2002)3663-3680, arXiv:hep-th/0202142.

[4] Richard Easther, Brian R. Greene, William H. Kinney, Gary Shiu, Inflation as a Probe of Short Distance Physics, Phys.Rev.D 64(2001)103502, arXiv:hep-th/0104102; Richard Easther, Brian R. Greene, William H. Kinney, Gary Shiu, Imprints of Short Distance Physics On Inflationary Cosmology, Phys.Rev.D 67(2003)063508, arXiv:hep-th/0110226; Richard Easther, William H Kinney, Hiranya Peiris, Observing Trans-Planckian Signatures in the Cosmic Microwave Background, JCAP 0505(2005)009, arXiv:astro-ph/0412613; Richard Easther, William H. Kinney, Hiranya Peiris, Boundary Effective Field Theory and Trans-Planckian Perturbations: Astrophysical Implications, JCAP 0508(2005)001, arXiv:astro-ph/0505426.

[5] Nemanja Kaloper, Matthew Kleban, Albion Lawrence, Stephen Shenker, Signatures of Short Distance Physics in the Cosmic Microwave Background, Phys.Rev.D 66(2002)123510, arXiv:hep-th/0201158; Nemanja Kaloper, Matthew Kleban, Albion Lawrence, Stephen Shenker, Leonard Susskind, Initial Conditions for Inflation, JHEP 0211(2002)037, arXiv:hep-th/0209231.

[6] Rachel Bean, Xingang Chen, Hiranya V. Peiris, Jiajun Xu, Comparing Infrared Dirac-Born-Infeld Brane Inflation to Observations, 0710.1812[hep-th]; Rachel Bean, Sarah E. Shandera, S. H. Henry Tye, Jiajun Xu, Comparing Brane Inflation to WMAP, JCAP 05(2007)004, arXiv:hep-th/0702107.

[7] R. Brandenberger and P. Ho,Noncommutative space-time, stringy space-time uncertainty principle, and density fluctuations, Phys.Rev.D 66(2002)023517, arXiv:hep-th/0203119; S. Alexander, R. Brandenberger, J. Magueijo, Noncommutative inflation, Phys.Rev.D 67(2003)081301, arXiv:hep-th/0108190;

[8] Qing-Guo Huang, Miao Li, CMB Power Spectrum from Noncommutative Spacetime, JHEP 0306(2003)014, arXiv:hep-th/0304203; Qing-Guo Huang, Miao Li, Noncommutative Inflation and the CMB Multipoles, JCAP 0311(2003)001, arXiv:astro-ph/0308458; Qing-Guo Huang, Miao Li, Running Spectral Index in Noncommutative Inflation and WMAP Three Year Results, Nucl.Phys.B 755(2006)286-294, arXiv:astro-ph/0603782; Qing-Guo Huang, Slow-roll reconstruction for running spectral index, JCAP 0611(2006)004, arXiv:astro-ph/0610924.

[9] X. Chen, M. Huang, S. Kachru and G. Shiu,Observational Signatures and Non-Gaussianities of General Single Field Inflation, JCAP 0701(2007)002, arXiv:hep-th/0605045.

[10] C. Armendariz-Picon, T. Damour and V. Mukhanov, k-inflation,Phys.Lett.B 458(1999)209, hep-th/9904075; J. Garriga and V. F. Mukhanov, Perturbations in k-inflation, Phys.Lett.B 458(1999)219-225, arXiv:hep-th/9904176.

[11] Kejie Fang, Bin Chen, Wei Xue, Non-commutative Geometry Modified Non-Gaussianities of Cosmological Perturbation, arXiv:0707.1970[hep-th].
[12] F. Lizzi, G. Mangano, G. Miele and M. Peloso, Cosmological perturbations and short distance physics from noncommutative geometry, JHEP 0206(2002)049, arXiv:hep-th/0203099.
[13] M. Alishahiha, E. Silverstein and D. Tong, DBI in the Sky, Phys.Rev.D 70(2004)123505, arXiv:hep-th/0404084.
[14] William H. Kinney, Konstantinos Tzirakis, Quantum modes in DBI inflation: exact solutions and constraints from vacuum selection, arXiv:0712.2043[astro-ph].
[15] Yi-Fu Cai, Yi Wang, Hawking-Moss Tunneling in Noncommutative Eternal Inflation, JCAP 0801:001,2008, arXiv:0711.4423; Yi-Fu Cai, Yi Wang, Is Noncommutative Eternal Inflation Possible? JCAP 0706:022,2007, arXiv:0706.0572; Yi-fu Cai, Yun-Song Piao, Probing noncommutativity with inflationary gravitational waves, Phys. Lett. B657:1-9,2007, arXiv:gr-qc/0701114.
[16] Richard Easther, Hiranya Peiris, Implications of a Running Spectral Index for Slow Roll Inflation, JCAP 0609(2006)010, arXiv:astro-ph/0604214.
[17] Running of the Scalar Spectral Index from Inflationary Models , Daniel J.H. Chung, Gary Shiu, Mark Trodden, Phys.Rev.D 68(2003)063501, arXiv:astro-ph/0305193.
[18] Hiranya Peiris, Richard Easther, Recovering the Inflationary Potential and Primordial Power Spectrum With a Slow Roll Prior: Methodology and Application to WMAP 3 Year Data, JCAP 0607(2006)002, arXiv:astro-ph/0603587; Hiranya V. Peiris , Daniel Baumann, Brett Friedman, Asantha Cooray, Phenomenology of D-Brane Inflation with General Speed of Sound, arXiv:0706.1240[astro-ph].
[19] Jie Ren, Hong-Guang Zhang,Xin-He Meng, Probing trans-Planckian physics and the curvature effect from primordial power spectrum with WMAP3 observations, arXiv:astro-ph/0612011; Xin-He Meng, Jie Ren, Hong-Guang Zhang, Exact solutions of embedding the 4D Universe in a 5D Einstein manifold, arXiv:arXiv:astro-ph/0701598.
[20] Yin-Zhe Ma, Xin Zhang, Brane inflation revisited after WMAP five year results, arXiv:0812.3421.
[21] G. Calcagni, Phys. Rev. D 70, 103525 (2004), arXiv:hep-th/0406006; G. Calcagni, Phys. Lett. B 606, 177 (2005), arXiv:hep-ph/0406057; G. Calcagni and S. Tsujikawa, Phys. Rev. D 70, 103514 (2004), arXiv:astro-ph/040754; G. Calcagni, J. Cosmol. Astropart. Phys. JCAP10(2005)009, arXiv:astro-ph/0411773.
[22] A. Ashoorioon, J.L. Hovdebo, Robert B. Mann, Nucl.Phys.B727:63-76,2005, arXiv: gr-qc/0504135.
[23] Daniel Baumann et al., CMBPol Mission Concept Study: Probing Inflation with CMB Polarization arXiv:0811.3919.